



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

From (4) and (6'), all the odd ordered derivatives of  $f(x)$  are zero when  $x = 0$ . By successive applications of (6') we have (since  $f''(0) = \pm 2c^2$ )

$$f^{IV}(0) = 2c^4, \quad f^{VI}(0) = \pm 2c^6, \quad \dots, \quad f^{(4n)}(0) = 2c^{4n}, \quad f^{(4n+2)}(0) = \pm 2c^{4n+2}.$$

Hence, by Maclaurin's theorem,

$$f(x) = 2 \pm 2c^2x^2/2! + 2c^4x^4/4! \pm 2c^6x^6/6! + \dots.$$

This gives either

$$f(x) = 2 \cdot \cosh cx, \quad \text{or} \quad f(x) = 2 \cdot \cos cx,$$

to which solutions  $f(x) = 0$  should be added as before.

Also solved by H. C. FEEMSTER, OSCAR S. ADAMS, PAUL CAPRON, and the PROPOSER.

#### MECHANICS.


##### 319. Proposed by LAENAS G. WELD, Pullman, Illinois.

A hexagonal pencil lies upon the inclined top of a drawing table and is on the point of either rolling or sliding. Find the angle between its direction and the horizontal edge of the table, the coefficient of friction being  $\mu$ .

#### SOLUTION BY H. S. UHLER, Yale University.

The condition for being on the verge of sliding is expressed by  $\mu = \tan \alpha$ , where  $\alpha$  denotes the angle between the table top and the horizontal and is sometimes called the "limiting angle of repose." The proof of this relation seems superfluous in this place because it is given in practically all text-books which are devoted in part or entirely to elementary mechanics.

In order that the pencil may be on the point of rolling, it is obviously necessary and sufficient that the vertical line through the center of gravity intersect the lateral edge which is at the lowest level. In the diagram let  $\overline{AB}$  and  $\overline{AC}$  indicate respectively a horizontal line on the table top and the lowest lateral edge of the hexagonal prism.  $G$  marks the center of gravity which is assumed to lie in the geometric axis of the pencil.  $\overline{GI}$  denotes the vertical through  $G$ , that is,



the line of action of the weight of the pencil which intersects the edge  $\overline{AC}$  in the point  $I$ .  $\overline{GE}$  and  $\overline{GF}$  are perpendiculars dropped from  $G$  upon the edge  $\overline{AC}$  and the table top, respectively. In other words, the plane  $EFG$  contains a right section of the prism passing through the center of gravity. Without quoting the well-known theorems of elementary geometry we see at once that  $\angle EFI = \beta$ ,  $\angle EGF = 30^\circ$ ,  $\angle FGI = \alpha$ ,  $\angle EFG = \angle FEI = \angle GFI = 90^\circ$ . Consequently,  $\overline{FG}/\overline{EF} = \cot 30^\circ = \sqrt{3}$ ,  $\overline{EF}/\overline{FI} = \cos \beta$ , and  $\overline{FI}/\overline{FG} = \tan \alpha$ . Multiplying these three equations together we obtain  $1 = \sqrt{3} \cos \beta \tan \alpha$ , but  $\tan \alpha = \mu$ , hence

$$\beta = \cos^{-1} \left( \frac{1}{\sqrt{3}\mu} \right).$$

*Remark.*—Since the greatest value of a cosine is unity the least value of  $\mu$  is equal to  $1/\sqrt{3} \doteq 0.57735$  corresponding to  $\alpha = 30^\circ$  and  $\beta = 0^\circ$ , as it should be.

Also solved by W. J. THOME, M. R. BOWERMAN, and GEORGE PAASWELL.